

Three Steps to Staging and Grading a Patient

Step 1: Initial Case Overview to Assess Disease

Screen:

- Full mouth probing depths
- Full mouth radiographs
- Missing teeth

Mild to moderate periodontitis will typically be either Stage I or Stage II

Severe to very severe periodontitis will typically be either Stage III or Stage IV

Step 2: Establish Stage

For mild to moderate periodontitis (typically Stage I or Stage II):

- Confirm clinical attachment loss (CAL)
- Rule out non-periodontitis causes of CAL (e.g., cervical restorations or caries, root fractures, CAL due to traumatic causes)
- Determine maximum CAL or radiographic bone loss (RBL)
- Confirm RBL patterns

For moderate to severe periodontitis (typically Stage III or Stage IV):

- Determine maximum CAL or RBL
- Confirm RBL patterns
- Assess tooth loss due to periodontitis
- Evaluate case complexity factors (e.g., severe CAL frequency, surgical challenges)

Step 3: Establish Grade

- Calculate RBL (% of root length x 100) divided by age
- Assess risk factors (e.g., smoking, diabetes)
- Measure response to scaling and root planing and plaque control
- Assess expected rate of bone loss
- Conduct detailed risk assessment
- Account for medical and systemic inflammatory considerations

Staging and Grading Periodontitis




The 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions resulted in a new classification of periodontitis characterized by a multidimensional staging and grading system. The charts below provide an overview. Please visit perio.org/2017wwdc for the complete suite of reviews, case definition papers, and consensus reports.

PERIODONTITIS: STAGING

Staging intends to classify the severity and extent of a patient's disease based on the measurable amount of destroyed and/or damaged tissue as a result of periodontitis and to assess the specific factors that may attribute to the complexity of long-term case management.

Initial stage should be determined using clinical attachment loss (CAL). If CAL is not available, radiographic bone loss (RBL) should be used. Tooth loss due to periodontitis may modify stage definition. One or more complexity factors may shift the stage to a higher level. See perio.org/2017wwdc for additional information.

	Periodontitis	Stage I	Stage II	Stage III	Stage IV
Severity	Interdental CAL <i>(at site of greatest loss)</i>	1 – 2 mm	3 – 4 mm	≥5 mm	≥5 mm
	RBL	Coronal third (<15%)	Coronal third (15% - 33%)	Extending to middle third of root and beyond	Extending to middle third of root and beyond
	Tooth loss <i>(due to periodontitis)</i>	No tooth loss		≤4 teeth	≥5 teeth
Complexity	Local	<ul style="list-style-type: none"> • Max. probing depth ≤4 mm • Mostly horizontal bone loss 	<ul style="list-style-type: none"> • Max. probing depth ≤5 mm • Mostly horizontal bone loss 	In addition to Stage II complexity: <ul style="list-style-type: none"> • Probing depths ≥6 mm • Vertical bone loss ≥3 mm • Furcation involvement Class II or III • Moderate ridge defects 	In addition to Stage III complexity: <ul style="list-style-type: none"> • Need for complex rehabilitation due to: <ul style="list-style-type: none"> – Masticatory dysfunction – Secondary occlusal trauma (tooth mobility degree ≥2) – Severe ridge defects – Bite collapse, drifting, flaring – < 20 remaining teeth (10 opposing pairs)
Extent and distribution	Add to stage as descriptor	For each stage, describe extent as: <ul style="list-style-type: none"> • Localized (<30% of teeth involved); • Generalized; or • Molar/incisor pattern 			



PERIODONTITIS: GRADING

Grading aims to indicate the rate of periodontitis progression, responsiveness to standard therapy, and potential impact on systemic health.

Clinicians should initially assume grade B disease and seek specific evidence to shift to grade A or C.

See perio.org/2017wwdc for additional information.

	Progression		Grade A: Slow rate	Grade B: Moderate rate	Grade C: Rapid rate
Primary criteria <i>Whenever available, direct evidence should be used.</i>	Direct evidence of progression	Radiographic bone loss or CAL	No loss over 5 years	<2 mm over 5 years	≥2 mm over 5 years
	Indirect evidence of progression	% bone loss / age	<0.25	0.25 to 1.0	>1.0
Case phenotype		Heavy biofilm deposits with low levels of destruction	Destruction commensurate with biofilm deposits	Destruction exceeds expectations given biofilm deposits; specific clinical patterns suggestive of periods of rapid progression and/or early onset disease	
Grade modifiers	Risk factors	Smoking	Non-smoker	<10 cigarettes/day	≥10 cigarettes/day
		Diabetes	Normoglycemic/no diagnosis of diabetes	HbA1c <7.0% in patients with diabetes	HbA1c ≥7.0% in patients with diabetes

The 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions was co-presented by the American Academy of Periodontology (AAP) and the European Federation of Periodontology (EFP).